

# $\gamma$ event anisotropy at RHIC

I.J. Johnson<sup>1,2</sup>, M. Kaneta<sup>2</sup>, and T.J. Symons<sup>2</sup> for the STAR collaboration  
<sup>1</sup>UC Davis, <sup>2</sup>LBNL

The first results on charged hadron event anisotropy in  $\sqrt{s_{NN}}=130$  GeV Au+Au collisions at RHIC were presented by the STAR experiment[1]. Additionally, anisotropies from identified hadrons ( $\pi^\pm$ ,  $K^\pm$ ,  $p$ , and  $\bar{p}$ ) have been presented in Ref.[2]. A question raised here is whether the event anisotropy for  $\pi^0$  is the same as for charged pions. The measurement of  $\gamma$  is necessary to measure  $\pi^0$  momenta since  $\pi^0 \rightarrow 2\gamma$  is the dominant decay mode. Here we would like to report progress in measuring  $v_2$  for  $\gamma$  rays as a first step of measurement of  $\pi^0 v_2$ .

We have established a method of gamma detection in the STAR experiment using the time projection chamber (TPC) to reconstruct electron positron pairs from photon conversions [3]. This method has the advantages that the momentum range ( $p_T > 100$  MeV) is wider and the energy resolution is better than typically the case for Electro-Magnetic Calorimeters (EMC). At the same time, the efficiency is quite low due to the small probability of gamma conversion ( $\approx 1\%$ ). These characteristics constrain our capability for gamma and  $\pi^0$  physics at low  $p_T$ .

Since the statistics of our data are limited, we have started to analyze  $v_2$  for gammas as a first step to measuring this quantity for  $\pi^0$ s. Figure. 1 shows gamma  $v_2$  as a function of transverse momentum. The  $v_2$  increases until about 2 GeV/c and then saturates. This feature is markedly similar to the charged hadron measurements[4]. Note that we need to take care of two corrections before we can extract the  $v_2$  of  $\pi^0$  from the gamma measurement; (1) Contributions from other gamma sources, e.g. direct gamma emission and hadron decays. (2) Kinematical effects which relate the gamma and  $\pi^0 v_2$  measurements.




Figure 1: STAR preliminary gamma  $v_2$  parameter (vertical) as a function of transverse momentum ( $p_T$ ) (horizontal), where  $v_2$  is second harmonic of azimuthal particle distribution. The error of  $v_2$  is statistical error only.

## References

- [1] K.H. Ackermann *et al.*, Phys. Rev. Lett.**86**, 402 (2001).

- [2] C. Adler *et al.*, Phys. Rev. Lett.**87**, 182301 (2001).  
 [3] I.J. Johnson, M. Kaneta, and T.J. Symons, 2000 NSD Annual Report.  
 [4] K.V.Filimonov *et al.*, 2001 NSD Annual Report.